

Coreceptor CD8-driven modulation of T cell antigen receptor specificity[☆]

Hugo A. van den Berg^{a,*}, Linda Wooldridge^b, Bruno Laugel^c, Andrew K. Sewell^b

^aWarwick Systems Biology Centre, Coventry House, University of Warwick, Coventry CV4 7AL, UK

^bDepartment of Medical Biochemistry and Immunology, University Hospital of Wales, Heath Park, Cardiff CF14 4XN, UK

^cLudwig Institute for Cancer Research, 155 Chemin des Boveresses, 1066 Epalinges, Suisse

Received 14 June 2007; received in revised form 25 July 2007; accepted 1 August 2007

Available online 8 August 2007

Abstract

The CD8 coreceptor modulates the interaction between the T cell antigen receptor (TCR) and peptide–major histocompatibility class I (pMHC I). We present evidence that CD8 not only modifies the affinity of cognate TCR/pMHC I binding by altering both the association rate and the dissociation rate of the TCR/pMHC I interaction, but modulates the sensitivity (triggering threshold) of the TCR as well, by recruiting TCR/pMHC I complexes to membrane microdomains at a rate which depends on the affinity of MHC I/CD8 binding. Mathematical analysis of these modulatory effects indicates that a T cell can alter its functional avidity for its agonists by regulating CD8 expression, and can rearrange the relative potencies of each of its potential agonists. Thus we propose that a T cell can specifically increase its functional avidity for one agonist, while decreasing its functional avidity for other potential ligands. This focussing mechanism means that TCR degeneracy is inherently dynamic, allowing each TCR clonotype to have a wide range of agonists while avoiding autorecognition. The functional diversity of the TCR repertoire would therefore be greatly augmented by coreceptor-mediated ligand focussing.

© 2007 Elsevier Ltd. All rights reserved.

Keywords: T cell immunity; T cell antigen receptor; CD8 coreceptor; Tetrameric major histocompatibility complex; Cross-reactivity; Polyspecificity

1. Introduction

Activation of cytotoxic T lymphocytes (CTLs) is dependent on recognition of protein antigens presented in the form of peptide fragments on the surface of target cells by class I products of the major histocompatibility complex (MHC I). The peptide–MHC I (pMHC I) complex interacts with the T cell antigen receptor (TCR), with an affinity governed by the TCR's complementarity-determining regions, which vary highly across the host's repertoire of TCR clonotypes (Malissen, 2003). The T cell surface glycoprotein CD8 independently interacts with invariable regions of the MHC I molecule (Wyer et al., 1999). Despite the low affinity of the pMHC I/CD8 interaction (Hutchinson et al., 2003),

TCR/pMHC I signalling is crucially dependent upon CD8 (Holler and Kranz, 2003), which acts as a coreceptor modulating the productivity of TCR engagement by pMHC I (Janeway, 1992; Luescher et al., 1995; Purbhoo et al., 2001) and thereby the TCR's functional avidity (Cawthon and Alexander-Miller, 2002).

Various distinct modulatory roles of CD8, possibly acting in concert, have been proposed: (i) promoting the association of TCR and pMHC I (Pecht and Gakamsky, 2005); (ii) stabilizing the TCR/pMHC I interaction (Luescher et al., 1995; Wooldridge et al., 2005), thus prolonging the mean dwell time of the interaction which alters the efficacy of the pMHC I ligand (Kalergis et al., 2001); and (iii) enhancing the rate at which the TCR/CD3 complex attains signalling status (Purbhoo et al., 2001; Holler and Kranz, 2003), by association of TCR/CD3 with protein tyrosine kinases such as p56^{lck} (Arcaro et al., 2001) and adaptor molecules such as LAT (Bosselut et al., 1999) and LIME (Brdičková et al., 2003). The contributions of

[☆]This work was supported by the Wellcome Trust.

*Corresponding author. Tel.: +44 24 76 52 3698;

fax: +44 24 76 52 4182.

E-mail address: hugo@maths.warwick.ac.uk (H.A. van den Berg).

these various coreceptor roles remain to be fully elucidated. Moreover, the CD8 $\alpha\beta$ heterodimer is considerably more potent as a coreceptor than the $\alpha\alpha$ homodimer (Bosselut et al., 2000; Arcaro et al., 2001; Gangadharan and Cheroutre, 2004), pointing to the importance of the third function, which is strongly dependent on the presence of the CD8 β chain (Bosselut et al., 2000; Arcaro et al., 2000).

In the present paper, we quantify the three CD8 coreceptor functions and evaluate their relative contributions to functional avidity, tying together a wealth of previous and novel data obtained in various experimental systems, using a mathematical theory of TCR avidity and T cell activation (reviewed in van den Berg and Rand, 2007). We examine TCR binding and functional avidity of both soluble and cell-surface expressed HLA A2 molecules that all have faithful interactions with the TCR yet exhibit a large range of CD8 binding affinities. In addition, we analyse data on a system where a suite of altered peptide ligands (APLs) is available with a range of affinities for a fixed CTL clone.

Our analysis supports the concept that CD8 is a key regulator of TCR degeneracy. Thus, we suggest that variation of CD8 expression allows the T cell to focus on the salient ligand by differentially adjusting its fine sensitivity to potential agonists (cf. Blok et al., 1996; Maile et al., 2005). Such focussing would allow each single T cell to have a wide range of potential agonists (Holler and Kranz, 2004; Wilson et al., 2004), while only one of these is a high-avidity ligand at any given time.

2. Materials and methods

Notation is summarized in Table 1.

2.1. Tetrameric pMHC1 staining, association and dissociation kinetics

Recombinant pMHC1 proteins were produced and multimerized as previously described (Wooldridge et al., 2005); gel filtration chromatography showed that >98.5% of the preparation was tetrameric. For tetramer association, 10^6 ILA1 CTLs were washed and resuspended in 150 μ l PBS with phycoerythrin-conjugated MHC1 tetramer at a final concentration of 1 μ g/ml (with respect to monomer). At the indicated time points, 10 μ l aliquots were taken of each sample and diluted to a final volume of 500 μ l in PBS prior to flow cytometric analysis. Tetramer concentrations were thus diluted 50-fold, ensuring that further staining after collection did not contribute significantly to the measured mean fluorescence intensity (MFI). Background staining was estimated by labelling ILA1 CTLs with non-cognate HLA A2 tetramers at room temperature for 30 min. This background value was subtracted from the MFI obtained at each time point. Detailed procedures for the tetramer decay assay have been described elsewhere (Wooldridge et al., 2005). Rate

Table 1
Glossary of notation

x_{ij}	surface density of i -plet TCR clusters j -fold bound
x	peptide incubation concentration
y	bioassay read out ($y_{min} \leq y \leq y_{max}$)
I	staining intensity
W	TCR signal (W_{bg} : background)
W_{act}	cellular activation threshold
w	TCR signal per pMHC1
Z	pMHC1 presentation level (\hat{Z} : maximum level)
R_i	surface density of i -plet TCR clusters
R_T	surface density of TCR molecules
ξ	saturation parameter
ω	location parameter (loglogistic distribution)
α	scale parameter (loglogistic distribution)
λ	empirical rate constant of tetramer association curve ($\lambda^{fast}, \lambda^{slow}, \lambda_{eff}$)
v	TCR/pMHC1 dissociation rate
v_i	i -plet TCR cluster decay rate
v^{slow}	TCR/pMHC1 dissociation rate when CD8 bound
v^{fast}	TCR/pMHC1 dissociation rate when not CD8 bound
μ	TCR/pMHC1 association rate in tetramer-induced cluster
δ	TCR loss rate from tetramer-induced cluster
γ	pMHC1/CD8 association rate
ρ	pMHC1/CD8 dissociation rate
ψ	pMHC1 tetramer recruitment rate
ϑ_1	TCR recruitment rate to singlet clusters
ϑ_2	TCR recruitment rate to duplet clusters
T_R	TCR receptor triggering threshold
T_R^o	TCR receptor triggering threshold in kinase-poor and/or phosphorylase-rich regions
T_R^*	TCR receptor triggering threshold in kinase-rich and/or phosphorylase-poor regions
T_{rc}	recruitment time to kinase-rich and/or phosphorylase-poor regions
ϑ_{rc}	recruitment rate to kinase-rich and/or phosphorylase-poor regions
K_{rc}	recruitment parameter
$G(\cdot)$	TCR receptor triggering probability function

parameters were estimated by non-linear least-squares regression.

2.2. Cells

Hmy.2 C1R B (C1R) cells expressing full length HLA A2 and mutants thereof were produced as previously described (Purbhoo et al., 2001). Cells were cloned and tested with anti-HLA A2 FITC (clone BB7.2, Serotec) to ensure that they expressed identical levels of MHC1 on their surface. The CTL clone NRT1 is specific for residues 77–85 (SLYNT-VAL) of HIV-1 p17 Gag, presented in association with HLA A2; NRT1 was generated from a healthy donor by tetramer sorting as previously described (Dunbar et al., 1998). The CTL clone ILA1 is specific for residues 540–548 (ILAKFLHWL) of the catalytic sub-unit of the ubiquitous tumour-associated telomerase reverse transcriptase (hTERT), presented in association with HLA A2; the monoclonal CTL line ILA1 was generated from a healthy donor by limiting dilution culture from a parent cell line

which was enriched for hTERT_{540–548}-specific T cells as previously described (Laugel et al., 2005).

2.3. Functional bioassays

CTLs were washed twice in RPMI/PSG, rested overnight in RPMI/PSG (with 2% FCS) and then adjusted to a concentration of 2.5×10^5 CTL/ml in RPMI/PSG (with 2% FCS). An aliquot of 10^6 C1R target cells was pulsed in a 50 μ l volume of various concentrations of SLYNTVATL peptide for 90 min at 37°C, 5% CO₂. After washing twice in RPMI/PSG, target cell concentration was adjusted to 2.5×10^5 cells/ml. An aliquot of 2.5×10^4 CTLs was placed into each well of a 96-U bottomed plate either with or without 2.5×10^4 pulsed C1R target cells in a final volume of 200 μ l. Plates were incubated for 4 h at 37°C. Supernatant was removed and assayed for MIP-1 β using a Quantikine ELISA kit (R&D systems), which is a very sensitive assay for CTL effector function.

2.4. Analysis of dose–response curves

Let y denote the assay read-out, approaching a maximum value y_{max} at high peptide pulsing concentrations, with a non-specific background value y_{min} . The ratio $(y - y_{min}) / (y_{max} - y_{min})$ can be taken to indicate the proportion of the T cells in the bioassay responding at the given pulsing concentration x . Accordingly, we assume that this proportion denotes the probability that a T cell will register a TCR-derived signal exceeding the cellular threshold for activation (Viola and Lanzavecchia, 1996; Hemmer et al., 1998; van den Berg and Rand, 2004a). This cellular threshold is denoted by W_{act} . We assume that the TCR signal W is composed of two terms, a background term W_{bg} due to signalling elicited by ligands other than the relevant agonist and a term due to the agonist, which we analyse as a product of the MHC I copy number (presentation level) Z and the agonist-derived TCR signal w , normalized per pMHC I molecule (van den Berg et al., 2001). It is standard to assume Langmuir binding for the presentation level: $Z(x) = \hat{Z}x / (\xi + x)$ where x denotes the peptide incubation concentration, ξ is a saturation parameter, and \hat{Z} is the maximum presentation level that can be attained. We assume that ξ is independent of the MHC I $\alpha 3$ mutation. Preliminary simultaneous least-squares fitting of this expression gave an estimate for ξ greater than the greatest pulsing concentration. Therefore we take $Z(x) = (\hat{Z} / \xi)x$.

We assume that $W_{act} - W_{bg}$ follows a lognormal distribution. For the sake of analytic convenience this can be approximated as a loglogistic distribution, with location and scale parameters $\omega > 0$ and $\alpha > 0$, respectively:

$$y = y_{min} + (y_{max} - y_{min})(1 + (x\hat{Z}(\omega\xi)^{-1}w)^{-\alpha})^{-1}. \quad (1)$$

The quantity $\hat{Z}(\omega\xi)^{-1}w$ can be estimated from the experimental data. This compound parameter serves as an index of TCR signal strength per ligand molecule (provided that

the quantity $\hat{Z}(\omega\xi)^{-1}$ has the same value for all ligands under consideration).

2.5. Theory of TCR triggering: TCR/pMHC I kinetics and optimal dwell time

We briefly review those aspects of the TCR/pMHC I kinetics and TCR triggering that are relevant for the quantification of coreceptor effects (for further mathematical details and arguments, see van den Berg et al., 2001, 2002; van den Berg and Rand, 2007; Burroughs and Wedagedera, 2006; Chan et al., 2003). The probability that an interaction between a pMHC I and a TCR molecule results in intracellular signalling is given by the following formula:

$$\mathbb{P}(\text{pMHC I triggers TCR}) = \int_0^\infty G(t) dF(t). \quad (2)$$

Here $G(t)$ denotes the probability that the TCR/CD3 complex and the associated (nascent) signalosome will have attained signalling status t time units after the TCR/pMHC I interaction started, and $F(t)$ is the distribution function of the lifetime of the TCR/pMHC I interaction, i.e. $F(t)$ denotes the probability that the TCR/pMHC I interaction will last for at most t time units. The standard assumption for the dwell-time distribution is that the interaction is characterized by a fixed hazard rate for dissociation, usually called the off-rate. This leads to the exponential distribution for the TCR/pMHC I dwell time: $F(t) = 1 - \exp(-vt)$, where v denotes the off-rate.

The function G reflects the steps involved in the transition to signalling status, the statistics of the weighting times associated with these steps, and their transition diagram (a graph expressing precedence relationships between these steps). It can be shown that the density G' is well approximated by a weighted series of Dirac pulses, provided that the transition diagram satisfies certain conditions on the number of steps relative to the graph's complexity. For the simplest member of this family of approximations, let $G(t)$ be the Heaviside step function with $G(t) = 0$ for $t \leq T_R$ and $G(t) = 1$ for $t > T_R$. Here T_R is a positive compound parameter which represents the time required for a TCR/pMHC I interaction to last in order to achieve intracellular signalling elicited by that TCR; T_R is called the *TCR receptor triggering threshold*.

On the standard assumption of a time-constant hazard rate v for TCR/pMHC I dissociation, the mean dwell time of the TCR/pMHC I interaction equals $1/v$ and the probability that the interaction will be productive works out as $\int_0^\infty G(t) dF(t) = \int_{T_R}^\infty dF(t) = \int_{T_R}^\infty \exp\{-vt\} dt/v = \exp\{-vT_R\}$. Under the assumption that the TCR/pMHC I interaction is *MHC-limited* (van den Berg et al., 2002), the rate at which a single pMHC I copy elicits intracellular signals is $v \exp\{-vT_R\}$. This is a hump-shaped function of v (Valitutti and Lanzavecchia, 1997; Bongrand and Malissen, 1998; Kalergis et al., 2001), with a maximum at $v = 1/T_R$, i.e. the optimum mean dwell time equals T_R .

Every TCR/pMHCI pair is characterized by a specific value for v . For a given TCR, virtually all pMHCI species are *null ligands* which dissociate very rapidly $v \gg T_R^{-1}$ whereas those ligands that satisfy $v \approx T_R^{-1}$ are precisely the agonists for that TCR clonotype (van den Berg et al., 2001; Lyons et al., 1996; Hudrisier et al., 1998; Matsui et al., 1994; van den Berg and Rand, 2004b).

A TCR may have one or several ligands such that $v \ll T_R^{-1}$ (Malissen, 2003); such rare *heteroclitic* agonists are suboptimal under MHC-limited conditions, but not under TCR-limited conditions (van den Berg et al., 2002). The mean rate at which a single agonist copy elicits TCR triggering events can be identified with the TCR's *functional avidity* for that ligand: a clonotype is said to have high avidity for a given epitope if it is able to respond even when the ligand is present at very low levels, whereas a low-avidity clone requires higher presentation levels for activation (Ashton-Rickardt and Tonegawa, 1994; Alexander-Miller, 2000; Gross et al., 2004).

2.6. Derivation of the microdomain recruitment equation

The coreceptor CD8 has been implicated in recruiting engaged TCR to specialized membrane microdomains that are kinase-enriched and/or phosphorylase-depleted. In such domains, phosphorylation and kinase/adaptor-recruitment processes are assumed to proceed more rapidly than outside these domains (Montixi et al., 1998; Janes et al., 1999; Arcaro et al., 2001; Doucey et al., 2001; Cawthon and Alexander-Miller, 2002; Bunnell et al., 2002; Filipp et al., 2003). These “favourable” domains may be dynamic regions of close approximation of the antigen-presenting cell (APC) and T cell membranes (Davis and van der Merwe, 2006).

Let T_R° denote the time it takes to attain a fully triggered TCR/CD3/signalosome outside of the microdomains, and let T_R^* denote the shorter corresponding time in these domains. Also, let T_{rc} denote the time it takes for an engaged TCR to be recruited to one of these domains; this is a random variable. The parameter T_R is a function of T_{rc} ; the shorter T_{rc} , the faster the TCR/CD3 proceeds through the steps that lead to activation. In particular, if T_{rc} is the time till recruitment, we have

$$T_R = \begin{cases} T_R^* + (T_R^\circ - T_R^*)T_{rc}/T_R^\circ & \text{for } T_{rc} \leq T_R^\circ, \\ T_R^\circ & \text{for } T_{rc} > T_R^\circ. \end{cases} \quad (3)$$

Let ϑ_{rc} denote the rate at which a TCR/pMHCI complex is recruited to a microdomain when bound to a CD8 capable of mediating the association with such a domain (Arcaro et al., 2000; Gangadharan and Cheroute, 2004; Gakamsky et al., 2005). The probability that a recruitment event occurs during an association with CD8 equals $\vartheta_{rc}/(\rho + \vartheta_{rc})$, where ρ denotes the pMHCI/CD8 dissociation rate. If pMHCI/CD8 kinetics is fast (Wyer et al., 1999) this probability is very small, which means that the expected time until recruitment can be treated as an

exponential variate with mean $(1 + \rho/\gamma)/\vartheta_{rc}$ where γ denotes the pMHCI/CD8 association rate (specifically, the moment generating function of T_{rc} is $M_{rc}(t) = (1 - \frac{\gamma+\rho}{p}t + \frac{\gamma\rho}{p}t^2)^{-1}$ where $p = \vartheta_{rc}/(\rho + \vartheta_{rc})$; this becomes the moment generating function of the exponential distribution upon dropping the term in t^2).

Conditioning the triggering probability on the recruitment time T_{rc} , we have

$$\begin{aligned} &\mathbb{P}(\text{pMHCI triggers TCR}) \\ &= \int_0^\infty \frac{\vartheta_{rc}\gamma e^{-\vartheta_{rc}\gamma T_{rc}/(\gamma+\rho)}}{\gamma + \rho} \int_0^\infty G(t; T_{rc}) dF(t) dT_{rc} \\ &= \frac{\vartheta_{rc}\gamma}{\gamma + \rho} \left(\int_0^{T_R^\circ} \exp\{-(\vartheta_{rc}\gamma T_{rc}/(\gamma + \rho) \right. \\ &\quad \left. + v(T_R^* + (T_R^\circ - T_R^*)T_{rc}/T_R^\circ))\} dT_{rc} \right. \\ &\quad \left. + \int_{T_R^\circ}^\infty \exp\{-(\vartheta_{rc}\gamma T_{rc}/(\gamma + \rho) + vT_R^\circ)\} dT_{rc} \right). \quad (4) \end{aligned}$$

The recruitment equation, which appears below as Eq. (19), is obtained from Eq. (4) with the definition $\kappa_{rc} = \vartheta_{rc}K_D^{pMHCI/CD8}\gamma/\rho$ and the physically reasonable assumptions $v \gg \vartheta_{rc}\gamma/(\gamma + \rho)$, $\rho/\gamma \gg 1$ and $\rho/\gamma \propto K_D^{pMHCI/CD8}$.

2.7. Kinetic theory of T cell/pMHCI-tetramer association and dissociation

Kinetic constants of the TCR/pMHCI interaction can be inferred from the intensity of fluorescence staining with pMHCI tetramers, as well as the rates at which a T cell acquires and loses the staining in association and dissociation assays. A mathematical model of these association and dissociation kinetics is depicted schematically in Fig. 1; the corresponding equations are given in Table 2. Steric considerations suggest that each tetramer can bind at most three TCR molecules. When one of the contacts




Fig. 1. Diagram showing the progressive binding of pMHCI tetramers to TCR singlets, duplets, and triplets. Non-covalent association and dissociation steps are indicated by arrows, which are labelled with the relevant rate constants.

Table 2
Equations of MHCI tetramer/T cell kinetics

$$\begin{aligned}\dot{x}_{11} &= \psi R_0 - (v + \vartheta_1 R_0)x_{11} \\ \dot{x}_{21} &= 2vx_{22} - (v + \vartheta_2 R_0 + 3\mu)x_{21} \\ \dot{x}_{22} &= \vartheta_1 R_0 x_{11} + 3\mu x_{21} - (2v + \vartheta_2 R_0)x_{22} \\ \dot{x}_{31} &= 2vx_{32} - (v + 6\mu)x_{31} \\ \dot{x}_{32} &= \vartheta_2 R_0 x_{21} + 6\mu x_{31} + 3vx_{33} - 2(v + \mu)x_{32} \\ \dot{x}_{33} &= \vartheta_2 R_0 x_{22} + 2\mu x_{32} - 3vx_{33}\end{aligned}$$

dissociates, the temporarily unoccupied MHCI site will generally be able to rebind the TCR. Consequently, the tetramer forms a comparatively persistent association with a cluster of TCRs. These associations can outlast the duration of the mean single-site dwell time by several orders of magnitude (Laugel et al., 2005); this *avidity* effect is most pronounced in *triplet* TCR clusters (defined by a 3:1 TCR/tetramer stoichiometry), somewhat less marked for *duplet* TCR clusters (2:1), and absent in TCR *singlets* (1:1).

As shown in Fig. 1, an MHCI tetramer newly recruited to the T cell surface forms a singlet which needs to recruit two more TCRs to form a triplet cluster. The parameters are: v , the single-site dissociation rate of the TCR/pMHCI interaction; μ , the single-site (re)binding rate within a TCR cluster (not necessarily identical to the native TCR/pMHCI association rate in a physiological cell:cell conjugate); δ , the rate of loss of an unbound TCR molecule from the cluster; and ψR_0 , the rate at which MHCI tetramers are recruited from the solution, where R_0 denotes the density of free TCR molecules on the T cell surface, i.e. TCRs which are not engaged in a tetramer-binding cluster. The ambient solution serves as an infinite reservoir, so that the concentration of MHCI tetramer in solution (to which ψ is proportional) can be taken to be constant. Moreover, $\vartheta_1 R_0$ denotes the rate at which singlet TCR clusters recruit another TCR molecule to form a duplet cluster and $\vartheta_2 R_0$ is the rate at which duplet TCR clusters recruit another TCR molecule to form a triplet cluster. The integer coefficients in Fig. 1 reflect the number of available sites in the various transitions.

When an MHCI tetramer is released to the ambient solution, the cluster will disband by diffusion, merging into the background of free TCRs. We assume that this break-up is effectively complete before one of the TCRs captures another tetramer from the solution. The only non-linearity in the system arises as a result of the following conservation law:

$$\begin{aligned}R_0 &= R_T - (R_1 + 2R_2 + 3R_3) \\ \text{with } R_1 &= x_{11}, \quad R_2 = x_{21} + x_{22}, \quad R_3 = x_{31} + x_{32} + x_{33}.\end{aligned}\tag{5}$$

where R_T is the surface density of all TCR molecules and x_{ij} denotes the surface density of TCR clusters containing i TCR molecules which are bound to the MHCI tetramer at j sites; we assume that $j = 4$ is ruled out by steric hindrance. Conjugated with a fluorescent group, the

MHCI tetramers can be used as a staining agent. Staining intensity I equals $\sum_{i=1}^3 \sum_{j=1}^i x_{ij}$ provided that tetramer fluorescent brightness is not affected by the valency of its coupling to TCR.

2.7.1. Derivation of the equilibrium staining equation

The following definitions are introduced for notational convenience:

$$v_1 = v, \quad v_2 = \frac{2v^2}{3\mu}, \quad v_3 = \frac{v^3}{2\mu^2}.\tag{6}$$

Assuming $\mu \gg v \gg \delta$ and $v_i \gg \vartheta_i R_T$, $i \in \{1, 2\}$, we find the following equations characterizing the steady state:

$$R_1 = \frac{\psi}{v_1} R_0, \quad R_2 = \frac{\psi \vartheta_1}{v_1 v_2} R_0, \quad R_3 = \frac{\psi \vartheta_1 \vartheta_2}{v_1 v_2 v_3} R_0,$$

which leads to a system of two equations describing staining intensity:

$$\begin{aligned}\frac{I}{R_T} &= \frac{\psi}{v_1} \left(\frac{R_0}{R_T}\right) + \frac{\psi \vartheta_1 R_T}{v_1 v_2} \left(\frac{R_0}{R_T}\right)^2 \\ &\quad + \frac{\psi \vartheta_1 R_T \vartheta_2 R_T}{v_1 v_2 v_3} \left(\frac{R_0}{R_T}\right)^3,\end{aligned}\tag{7}$$

$$1 = \frac{\psi}{v_1} \left(\frac{R_0}{R_T}\right) + 2 \frac{\psi \vartheta_1 R_T}{v_1 v_2} \left(\frac{R_0}{R_T}\right)^2 + 3 \frac{\psi \vartheta_1 R_T \vartheta_2 R_T}{v_1 v_2 v_3} \left(\frac{R_0}{R_T}\right)^3\tag{8}$$

(the second equation is just the conservation law, Eq. (5)). Routine analysis yields $I \rightarrow R_T/3$ as $v/\mu \rightarrow 0$, which shows that most of the staining involves TCR triplets when rebinding is rapid. Moreover, if we assume that v is proportional to the off-rate of the TCR/pMHCI interaction and that ψ , ϑ_1 , ϑ_2 , and μ are all proportional to the on-rate of this interaction, we can rewrite these equations in terms of the dissociation constant $K_D^{TCR/pMHCI}$. In particular, the staining equation becomes

$$\frac{I}{R_T} = \frac{K_A}{K_D} \cdot \frac{R_0}{R_T} + \left(\frac{K_B}{K_D}\right)^3 \left(\frac{R_0}{R_T}\right)^2 + \left(\frac{K_C}{K_D}\right)^6 \left(\frac{R_0}{R_T}\right)^3,\tag{9}$$

where $K_D = K_D^{TCR/pMHCI}$ and the constants K_A , K_B , and K_C absorb the various proportionality constants. The sixth-power dependence of the affinity of the TCR/pMHCI interaction is the result of two avidity interactions: one involving the dominance of triply-bound tetramers, as discussed above, and one involving the progression of binding through TCR singlets, duplets, and triplets. These results imply a sigmoid dependence of I on $K_D^{TCR/pMHCI}$. In particular, rapid within-cluster rebinding makes the sixth-power term dominant, so that this curve becomes very steep: the transition from background staining to saturated staining is predicted to occur within less than an order of magnitude, as was confirmed by experimental observations (Laugel et al., 2007).

2.7.2. Effective dissociation rate of the tetramer/TCR-triplet complex

In a tetramer dissociation assay, tetramer-treated T cells are placed in a tetramer-free solution which contains antibodies to cap tetramers coming off the T cell surface, precluding their re-attachment. If the rebinding rate μ far exceeds the single-site dissociation rate ν and most of the MHC I tetramer bound to the surface is bound to TCR triplet clusters, staining intensity decays with apparent rate constant ν_3 . The kinetics of the triplet system $\{x_{31}, x_{32}, x_{33}\}$ is linear and thus there are two further eigenvalues. A routine exercise in analysis yields the following (approximate) formulae for these two additional eigenvalues:

$$-6\mu - 4\nu \quad \text{and} \quad -2(\mu + \nu).$$

These additional eigenvalues correspond to equilibration between singly, double, and triply bound tetramers in the triplet cluster. Inasmuch as these two rates exceed ν_3 (since $\mu \gg \nu$) by several orders of magnitude, the empirical dissociation curve (as observed on the ν_3^{-1} time scale) will be effectively monophasic. The observed rate constant $\lambda_{\text{eff,off}}$ then serves as an estimate for ν_3 .

Inasmuch as μ is unknown, the single-site off-rate ν cannot be determined directly from the measured rate $\lambda_{\text{eff,off}}$ (i.e. ν_3). However, *ratios* of the experimental rates (e.g. between mutant variants) can be used to gauge the effect on ν ; since $\nu_3 \propto \nu^3$ the cube root of the experimental ratio gives the relative change of ν . When initial staining is predominantly in the form of duplets, a similar argument applies, but now the rate ν_2 is experimentally gauged and a square root has to be extracted.

2.7.3. Analysis of MHC I-tetramer/T cell association kinetics

The association phase of tetramer staining has more complicated dynamics than the decay curves, since it involves a progression through the binding states depicted in Fig. 1. When $\mu \gg \nu$ (rapid within-cluster rebinding), fast modes within the triplet and duplet systems rapidly relax the dynamics onto the $\{R_1, R_2, R_3\}$ -manifold, on which the following dynamics obtains:

$$\dot{R}_1 = \psi R_0 - (\nu_1 + \vartheta_1 R_0) R_1, \quad (10)$$

$$\dot{R}_2 = \vartheta_1 R_0 R_1 - (\nu_2 + \vartheta_2 R_0) R_2, \quad (11)$$

$$\dot{R}_3 = \vartheta_2 R_0 R_2 - \nu_3 + R_3, \quad (12)$$

which immediately yields the following staining kinetics:

$$\dot{I} = \psi R_T - \lambda_{\text{eff}}(t) I \quad (13)$$

with a *time-varying* effective rate $\lambda_{\text{eff}}(t)$, defined by

$$\lambda_{\text{eff}}(t) = \sum_{i=1}^3 \frac{(i\psi + \nu_i) R_i(t)}{R_T - R_0(t)}. \quad (14)$$

Eq. (13) shows that $\dot{I}|_{t=0} = \psi R_T$; this means that the rate of MHC I tetramer recruitment from solution can be estimated from the initial slope of the association curve.

Furthermore, the equilibrium staining equation (9) can be used to calculate the steady-state value of λ_{eff} , because the relative magnitudes of the three terms on the right-hand side of Eq. (9) correspond directly to $R_i/(R_T - R_0)$, the relative numbers of singlets ($i = 1$), duplets ($i = 2$), and triplets ($i = 3$) at steady state.

3. Results

The TCR triggering rate per agonist molecule is given by the formula,

$$\nu \exp\{-\nu T_R\}$$

(van den Berg et al., 2002), which is illustrated in Fig. 2, where ν is the TCR/pMHC I dissociation rate and T_R is the time the TCR/pMHC I docking needs to last in order to promote intercellular signalling. At least two distinct coreceptor effects are suggested by the formula: modulation of the TCR/pMHC I dissociation rate ν (Hutchinson et al., 2003; Wooldridge et al., 2005) and modulation of the receptor triggering threshold time T_R by colocalization with kinases, adaptors, and linkers (Doucey et al., 2001; Montixi et al., 1998; Bosselut et al., 1999; Brdičková et al., 2003; Filipp et al., 2003). We investigate both types of coreceptor-mediated modulations.




Fig. 2. Graph illustrating the scope for coreceptor-mediated modulation of the TCR activation rate per pMHC I molecule. A reduction of the TCR/pMHC I dissociation rate ν by 50% takes the ligand at point a to b. However, the same ligand is taken to point c by a 50% reduction in the receptor threshold T_R . The two modulatory effects can act synergistically: when both ν and T_R are reduced by 50%, the ligand at point a is taken to d. In contrast, the optimal ligand at point e becomes less effective when the TCR/pMHC I dissociation rate is reduced by 50% (e to f) whereas the same ligand is improved by a 50% reduction in the receptor threshold T_R (e to g).

3.1. Coreceptor modulation of the mean dwell time of the TCR/pMHC interaction

One important effect of the MHC/CD8 interaction is a reduction of v , the TCR/pMHC dissociation rate (Table 3; Luescher et al., 1995; Wooldridge et al., 2005). Thus, let v^{slow} denote the dissociation rate when MHC is bound to CD8, and v^{fast} otherwise. Coreceptor-mediated stabilization of the TCR/pMHC interaction is then expressed by the inequality $v^{fast} > v^{slow}$. A useful index for the stabilizing effect is v^{slow}/v^{fast} ; this ratio is at least 2 (Wooldridge et al., 2005; Laugel et al., 2007). To relate this index to T cell activation, consider first the distribution function $F(t)$. In the presence of CD8, $F(t)$ is given as $1 - (S^{fast}(t) + S^{slow}(t))$ where the latter two terms satisfy the differential equations,

$$\dot{S}^{fast}(t) = \rho S^{slow}(t) - (v^{fast} + \gamma)S^{fast}(t), \tag{15}$$

$$\dot{S}^{slow}(t) = \gamma S^{fast}(t) - (v^{slow} + \rho)S^{slow}(t) \tag{16}$$

with initial conditions $S^{fast}(0) = \rho/(\gamma + \rho)$ and $S^{slow}(0) = \gamma/(\gamma + \rho)$. Here γ is the rate of MHC association with CD8, which is proportional to the density of free CD8 molecules on the T cell surface ($[CD8]_F$) and ρ is the MHC/CD8 dissociation rate. Solving Eqs. (15) and (16), we obtain expressions for $F(t)$, best understood by

consideration of two special cases representing the extremes of a range of possibilities.

First, in the case where pMHC/CD8 kinetics is very *rapid* compared to the TCR/pMHC interaction (i.e. $\min(\gamma, \rho) \gg v^{fast}$) we obtain

$$F(t) = 1 - \exp\{(\gamma + \rho)^{-1}(\rho v^{fast} + \gamma v^{slow})t\}. \tag{17}$$

The mean dwell time of the TCR/pMHC interaction is then found as

$$\int_0^\infty t dF(t) = (\gamma + \rho)/(\rho v^{fast} + \gamma v^{slow}).$$

On the other hand, when pMHC/CD8 kinetics is very *slow* compared to the cognate interaction (i.e. $\max(\gamma, \rho) \ll v^{slow}$), so that a TCR/pMHC interaction typically occurs in its entirety with the coreceptor either associated or disassociated, we have

$$F(t) = 1 - \frac{\rho}{\gamma + \rho} \exp\{-v^{fast}t\} - \frac{\gamma}{\gamma + \rho} \exp\{-v^{slow}t\} \tag{18}$$

and the mean dwell-time of the TCR/pMHC interaction is

$$\int_0^\infty t dF(t) = (\rho/v^{fast} + \gamma/v^{slow})/(\gamma + \rho).$$

To illustrate how the T cell can differentially modulate its sensitivity to various ligands by altering its surface expression of CD8, the rate of TCR triggering is plotted as a function of the scaled free CD8 density in Fig. 3. The TCR triggering rate is expressed by the probability of productive TCR/pMHC interaction, divided by mean dwell time. The curves obtain for hypothetical pMHC ligands which differ with respect to their TCR/pMHC dissociation rates (the effect magnitude v^{slow}/v^{fast} is assumed the same for all ligands, for the sake of simplicity). Scaling is with respect to the two-dimensional dissociation constant of the pMHC/CD8 interaction. Accordingly, the scaled free CD8 density is the net representation not only of the cell surface expression level of CD8, but also of any

Table 3
MHC tetramer staining: dissociation rates

Ligand	Dissociation rate		Increase (%)	
	Wild-type	CD8-null	Triplets	Duplets
3G	0.123	0.462	55	92
8Y	0.203	0.587	42	69
8T	0.206	0.608	43	72

Dissociation rate expressed in min^{-1} . Wild-type: HLA-A2; CD8-null: HLA-A2 DT227/8KA; increase: calculated increase of off-rate in CD8-null.




Fig. 3. Normalized (maximum = 1) rate of TCR triggering dependent on the free CD8 density on the T cell surface (normalized with respect to the dissociation constant) for various ligands, in the case of fast CD8 kinetics, $\min(\gamma, \rho) \gg v^{fast}$ (A) or slow CD8 kinetics, $\max(\gamma, \rho) \ll v^{slow}$ (B). For all ligands, the magnitude of the CD8 stabilization effect has been set at $v^{slow}/v^{fast} = 20$ in these graphs, in order to clearly demonstrate the effects. LOW CD8 denotes a hypothetical ligand that is most potent at low CD8 densities, and similarly for INTERMEDIATE CD8 and HIGH CD8.

physico-chemical properties of CD8 that affect the association and dissociation rates of the pMHC/CD8 interaction.

As the scaled free CD8 density increases, the TCR/pMHC dissociation rate changes over from v^{fast} to v^{slow} . Along with this shift, the potency of a ligand shifts as well: a ligand such that $v^{fast} \approx T_R^{-1}$ is more potent at low CD8 and becomes less potent as the free CD8 density increases, whereas the opposite is true for a ligand for which $v^{slow} \approx T_R^{-1}$. These effects are illustrated by the CD8–response curves in Fig. 3. Variation of CD8 surface expression thus equips the T cell with a tuning mechanism that allows a given TCR clonotype to control ligand promiscuity by optimizing responsiveness to the relevant ligand and at the same time reducing responsiveness to other potential agonists (which would be favoured at different CD8 expression levels). In effect, the coreceptor CD8 mediates focussing on a specific ligand chosen from among a larger set of potential agonists.

Such CD8-mediated focussing may also be involved in the maturation of functional responsiveness during an immune response, irrespective of additional maturation mechanisms such as clonal selection or TCR editing. The scope for tuning is somewhat greater when pMHC/CD8 kinetics are rapid compared to the mean TCR/pMHC

dwell time (Fig. 3A), where a tuning to intermediate ligands is possible. This may indeed be the case as the available evidence (Wyer et al., 1999) suggests that pMHC/CD8 kinetics is relatively rapid.

3.2. Coreceptor modulation of the threshold duration of the TCR/pMHC interaction

To assess a further CD8 enhancement effect besides dwell-time prolongation, functional response bio-assays were carried out in which CTLs were incubated with APCs incubated with agonist peptide at increasing concentrations, and expressing various HLA mutants with differing CD8 affinities (Hutchinson et al., 2003; Wooldridge et al., 2005). The response of CTL clones stimulated by APCs incubated with the agonist HIV-1 Gag epitope SLYNT-VATL, measured as the concentration of MIP-1 β produced after a 4 h incubation, is shown in Fig. 4A as a function of agonist peptide concentration. To compare the relative strengths of the MHC-specific TCR triggering rate elicited by these mutants, the MHC-specific triggering rates estimated from the dose–response curves were normalized with respect to the TCR triggering rate associated with mutant DT227/8KA; the HLA A2 heavy chain $\alpha 3$ domain of the latter mutant is incapable of




Fig. 4. MIP-1 β release response of NRT1 CTLs as a function of agonist incubation concentration. Response is shown (A) for APCs expressing various mutations in HLA A2: \circ : wild type; \square : Q115E ($\alpha 2$ domain); \triangle : A245V ($\alpha 3$ domain); \diamond : DT227/8KA ($\alpha 3$ domain). Curves represent the least-squares fit of Eq. (1), where $y_{min} = 7.9 \pm 0.07$ pg/ml; $y_{max} - y_{min} = 312 \pm 0.17$ pg/ml; $\alpha = 1.03 \pm 0.05$; $\log(\tilde{Z}(\omega\zeta)^{-1}w_{DT227/8KA}M^{-1}) = 5.10 \pm 0.03$; $\log(w_{Q115E}/w_{DT227/8KA}) = 3.58 \pm 0.06$; $\log(w_{wild\ type}/w_{DT227/8KA}) = 3.19 \pm 0.05$; $\log(w_{A245V}/w_{DT227/8KA}) = 2.52 \pm 0.07$. Values of $\log(w./w_{DT227/8KA})$ are plotted against the cube root of the effective off-rate in tetramer decay curves (B) and the dissociation constant of the pMHC/cb8 interaction (C). Cube roots of effective off-rates are as follows: Q115E: $0.42 \text{ min}^{-1/3}$; wild type: $0.44 \text{ min}^{-1/3}$; A245E: $0.54 \text{ min}^{-1/3}$; DT227/8KA: $0.85 \text{ min}^{-1/3}$. $K_D^{pMHC/cb8}$ (Q115E): $85 \mu\text{M}$; $K_D^{pMHC/cb8}$ (wild type): $128 \mu\text{M}$; $K_D^{pMHC/cb8}$ (A245V): $500 \mu\text{M}$ (data from Wooldridge et al., 2005).

binding to CD8 (Purbhoo et al., 2001). The relative strength of the MHCI-specific TCR triggering rate, thus defined, is shown as a function of the cube root of the effective tetramer decay rate in Fig. 4B, and as a function of $K_D^{pMHCI/CD8}$ in Fig. 4C. The cube root of the effective tetramer decay rate is proportional to the single-site TCR/pMHCI dissociation rate (see Wooldridge et al., 2005 and kinetic theory in Materials and methods).

The trends shown in Fig. 4B and C suggest an involvement of CD8 in the triggering of TCRs, in addition to the dwell-time effect discussed in the previous section. This second coreceptor mode of action would involve interaction with the molecular machinery which assembles the signalosome. The TCR triggering theory represents such modulation via the function G in Eq. (2), here taken to have the simplest possible form, a Heaviside function with threshold parameter T_R . In particular, this putative additional role of CD8 is detailed by the following assumptions: (i) pMHCI/CD8 kinetics is fast relative to TCR/pMHCI kinetics (Wyer et al., 1999; Gakamsky et al., 2005); (ii) the T cell membrane is spatially heterogeneous, with microdomains that favour TCR triggering, being rich in the required kinases (or poor in phosphatases, or both), versus unfavourable microdomains that are kinase-depleted and/or phosphatase-enriched (Bunnell et al., 2002; Davis and van der Merwe, 2006); (iii) TCRs can be recruited to the favourable domains through interaction with the coreceptor, dependent on the affinity of the CD8:α3 interaction (Doucey et al., 2001; Montixi et al., 1998); (iv) the rate of recruitment is slow relative to the mean duration of the pMHCI/CD8 contact; and (v) when the TCR/pMHCI ternary complex has been recruited to these membranes, it remains in the favourable domain for at least the duration of the TCR/pMHCI docking.

Letting w denote the MHC-specific TCR triggering rate, T_R° the triggering threshold in the unfavourable domains, T_R^* the threshold in the favourable domains ($T_R^\circ > T_R^*$), and κ_{rc} a recruitment parameter proportional to the surface density of CD8 molecules associated with the favourable domains, we obtain the following equation from assumptions (i)–(v):

$$w = v \exp\{-vT_R^\circ\} + \frac{\kappa_{rc}(T_R^\circ - T_R^*)/T_R^\circ}{K_D^{pMHCI/CD8}} \exp\{-vT_R^*\}. \quad (19)$$

The rate at which ternary complexes are recruited to the favourable domains is $\kappa_{rc}/K_D^{pMHCI/CD8}$ (one may take the two-dimensional dissociation constant to be proportional to the three-dimensional dissociation constant, for which data are available; the proportionality constant, known as the *confinement length*, is absorbed in the recruitment parameter). For the TCR:pMHCI dissociation rate v we have $v = C \sqrt[3]{k_{eff\ off}}$ where $k_{eff\ off}$ is the effective dissociation rate as measured in tetramer dissociation experiments (Wooldridge et al., 2005) and C is a correction factor with the dimensions $\text{TIME}^{-2/3}$. Given the estimates on $w_{\square}/w_{DT227/8KA}$ for $\square = Q115E$, wild type, or A245V,

Eq. (19) furnishes three equations in three unknown parameters, one of which is T_R°/T_R^* which is how much faster the steps in CD3 activation and signalosome assembly proceed in the favourable domains (steps in signalosome assembly include ITAM phosphorylation and recruitment of kinases and adaptors). The other two parameters are numerical conversion coefficients. The best solution of these equations by least-squares yields the estimate $T_R^\circ/T_R^* \approx 4.49$. While there is considerable uncertainty associated with this estimate, it does indicate that CD3 activation and kinase/adaptor recruitment proceeds faster in the favourable domains.

Since the T_R°/T_R^* factor ~ 4.5 occurs in the exponent (see Eq. (19)), its impact on the TCR triggering rate can be huge: for a pMHCI ligand with mean dwell time $\sim T_R^*$, the probability that the interaction is productive will be some 30 times greater when the ternary complex is recruited to the favourable domains as compared to the unfavourable domains. On the other hand, for a ligand optimal for the unfavourable domains (i.e. $v^{-1} \approx T_R^\circ$), the TCR triggering rate will be about twice as large in the unfavourable domains as compared to the favourable domains. Thus, the effect of dwell-time modulation described in the previous section will be strongly magnified whenever the contribution of favourable domains to the overall TCR triggering rate is significant. The enhanced TCR triggering rate is represented by the second term on the right-hand side of Eq. (19) which is proportional to the surface density of CD8 molecules associated with the favourable domains. In the absence of favourable domain-associated CD8 molecules, optimal ligands are those pMHCI with mean dwell time $v^{-1} \sim T_R^\circ$ whereas at high levels of such CD8s, ligands are optimal if their mean dwell time is near T_R^* .

Thus membrane heterogeneity plus CD8-mediated recruitment provides a second focussing mechanism that allows the clonotype to tune its sensitivity specifically to one pMHCI species from among a range of potentially optimal ligands.

3.3. Coreceptor modulation of TCR/pMHCI affinity

A series of hTERT_{540–548} altered peptide ligands (APLs) to CTL clone ILA1 was used to obtain estimates for the parameters of the staining equation (9) (see also Laugel et al., 2007). Singlet TCRs were found to contribute very

Table 4
MHCI tetramer staining: TCR triplet clusters

Ligand	Triplet fraction		$K_D^{TCR/pMHCI}$ (μM)
	Wild-type	CD8-null	
3G	0.993	0.984	3.7 ± 0.28
8Y	0.921	0.733	22.6 ± 2.05
8T	0.882	0.536	27.6 ± 4.71

Wild-type: HLA-A2; CD8-null: HLA-A2 DT227/8KA. K_D for the TCR/pMHCI interaction.

little to equilibrium staining, and K_A had to be fixed at the value zero; estimated values were $K_B = 10.8 \pm 1.10 \mu\text{M}$ and $K_C = 26.6 \pm 1.03 \mu\text{M}$. These values apply for the wild-type HLA A2 molecule; for interactions with HLA A2 DT227/8KA, these values have to be divided by a CD8 effect factor estimated at 2.25 ± 0.086 . Table 4 shows the proportion of staining clusters that are triplets rather than duplets, calculated from the staining equation, both for wild-type HLA and DT227/8KA, in three APLs, 3G, 8Y, and 8T. The TCR/pMHCI dissociation constants are listed for comparison; the trend is clearly for lower-affinity APLs to have an increasing proportion of clusters present as duplets; and this trend is emphasized when the pMHCI/CD8 interaction is abrogated.

The estimated CD8 affinity effect factor ~ 2.25 absorbs mean dwell-time prolongation (i.e. a decrease of the off-rate v mediated by CD8) as well as a possible effect on the on-rate. Consider the apparent MHC tetramer staining dissociation rates for wild-type HLA and HLA DT227/8KA, given in Table 3. The column marked “triplets” shows the estimated percentage increase of the single-site off-rate consequent upon abrogation of the pMHCI/CD8 interaction, based on all-triplet staining (extracting a cube root of the rate ratios) whereas the column marked “duplets” shows the percentages on the basis of all-duplet staining (based on the square root of the rate ratios). The effective dwell-time prolongation thus appears to be about 1.5, indicating that the effect of pMHCI/CD8 on the single-site on-rate is again about 1.5, to make up the total factor of 2.25. Thus, the data suggest that the favourable effect conferred by the coreceptor CD8 is composed of roughly equal contributions of increased on-rate and decreased off-rate.

Further evidence pointing to an on-rate modulatory effect of CD8 is provided by the staining association kinetics of the hTERT_{540–548} APLs system. The association curves (shown in Laugel et al., 2007) closely fit a biphasic exponential model:

$$I(t) = I_{max}^{fast}(1 - \exp\{-\lambda^{fast}t\}) + I_{max}^{slow}(1 - \exp\{-\lambda^{slow}t\}). \quad (20)$$

The estimated values for the empirical rate parameters are listed in Table 5. Numerical simulations of the full kinetic system (Table 2) confirm that such biphasic behaviour is consistent with the much more intricate underlying dynamics. As indicated in Materials and methods, the

initial rate of association is a measure of ψR_T , where the parameter ψ absorbs the incubation concentration of MHC tetramer as well as the rate at which singlet TCRs capture tetramers from solution. For the empirical model, Eq. (20), the initial rate is given by $I_{max}^{fast}\lambda^{fast} + I_{max}^{slow}\lambda^{slow}$. Combining these quantities for both wild-type HLA and HLA DT227/8KA, we arrive at an estimated percentage decrease of the single-site tetramer capture rate consequent upon abrogation of the pMHCI/CD8 interaction, shown in the column marked “capture effect” in Table 5. The effects are dramatic, and become more pronounced with lower TCR/pMHCI affinity.

The empirical data fits also allow the estimation of the time-varying effective rate of staining kinetics, $\lambda_{eff}(t)$; Eq. (13). This rate relaxes to the equilibrium value $\lambda_{eff}(\infty)$ on a time scale $\sim 1/\lambda^{fast}$. The equilibrium value can be calculated from the empirical parameters, as follows:

$$\lambda_{eff}(\infty) = \frac{I_{max}^{fast}\lambda^{fast} + I_{max}^{slow}\lambda^{slow}}{I_{max}^{fast} + I_{max}^{slow}} \quad (21)$$

(this follows immediately from Eq. (13)). These estimates, shown in the two final columns of Table 5, indicate that abrogation of the pMHCI/CD8 interaction leads to a decrease in $\lambda_{eff}(\infty)$, which is most pronounced in ligand 3G.

4. Discussion

This study attempts to dissect and quantitate the various ways in which CD8 modulates and augments TCR signal transduction. Our analysis suggests that, in addition to its role in recruiting the kinase p56^{lck} to the TCR/CD3 complex, which is essential for virtually all normal syngeneic interactions (Arcaro et al., 2001; Holler and Kranz, 2003; Lyons et al., 2006), the coreceptor CD8 has a number of modulatory effects: CD8 (i) enhances the TCR/pMHCI association rate by 50% or more; (ii) stabilizes the TCR/pMHCI interaction by at least 50%; and (iii) recruits TCR/pMHCI complexes to membrane domains where the steps needed to attain signalling status for the TCR/CD3 complex proceed more than four times faster, equivalent to a 30-fold increase in productive ligand engagement. These modulatory effects allow the T cell to fine-tune its sensitivity to the salient ligands.

Coreceptor modulation of functional avidity is differential: as illustrated in Figs. 2 and 3, the T cell becomes

Table 5
MHC tetramer staining: association rates

Ligand	λ^{slow} (min ⁻¹)		λ^{fast} (min ⁻¹)		Capture effect (%)	$\lambda_{eff}(\infty)$ (min ⁻¹)	
	Wild-type	CD8-null	Wild-type	CD8-null		Wild-type	CD8-null
3G	0.18	0.07	3.08	4.01	61	1.53	0.76
8Y	0.12	0.13	3.07	4.45	85	1.30	1.25
8T	0.14	0.13	8.10	7.81	90	3.11	2.34

Wild-type: HLA-A2; CD8-null: HLA-A2 DT227/8KA. Capture effect: calculated decrease of recruitment rate ψ in CD8-null. Data from Laugel et al. (2007).

more responsive to one of its potential agonists while becoming less responsive to another one at the same time. This differential modulation enables genuine focussing, as opposed to merely tuning signal transduction gain, which affects signals from all potential agonists equally (Slifka and Whitton, 2001; Schade and Levine, 2003). On this view, T cells exert active control over which antigen they focus on, by regulating the modulatory actions of the coreceptor in three ways: (i) through the expression level of CD8 (Blok et al., 1996; Maile et al., 2005); (ii) through the ratio of expression between CD8 $\alpha\alpha$ and CD8 $\alpha\beta$ (Cawthon and Alexander-Miller, 2002); and (iii) through the affinity of pMHC I for CD8, which can be modulated by varying the glycosylation of the interacting molecules (Daniels et al., 2001). Manipulation of one or more of these three control parameters should make it possible to re-order the hierarchy of potency among a CTL's potential agonists in an experimental system; if successful, such experiments would corroborate a central role for coreceptor-mediated fine tuning and focussing of TCR specificity against a background of high degeneracy.

The existence of a focussing mechanism implies that an individual clonotype may have a fairly wide range of ligands, among which, at any given time, only a small subset is potent (i.e. can act as optimal agonist). We thus propose that CD8-modulation of TCR promiscuity endows the TCR repertoire with an additional dimension of diversity, enhancing the capability of a limited number of clonotypes to cover "epitope space" efficiently (Goldrath and Bevan, 1999; Mason, 2001; Nicholson et al., 2000; Holler et al., 2002).

The inherent polyspecificity and cross-reactivity of the TCR has been well established (Gavin and Bevan, 1995; Ignatowicz et al., 1996; Kersh and Allen, 1996; Mason, 1998; Holler et al., 2002; Holler and Kranz, 2004; Bankovich et al., 2004; Wilson et al., 2004). Yet such TCR promiscuity needs to be reconciled with the need to distinguish salient antigens (pathogen-related non-self and harmful self) from non-salient ones (harmless self), and the immune system employs various central and peripheral tolerance mechanisms acting in unison (Seddon and Mason, 2000; Anderton and Wraith, 2002; van den Berg and Rand, 2004b; van den Berg and Molina-Paris, 2003).

Coreceptor-mediated focussing of TCR degeneracy may be an important additional mechanism. In particular, the concept of focussed modulation of functional avidity strongly complements theories that view degeneracy as a fundamental molecular feature of the TCR, such as the *multiple conformer* theory which proposes that a single TCR exists in multiple conformations that are in equilibrium (Holler and Kranz, 2004), allowing a single T cell to potentially recognize many different ligands. Moreover, coreceptor-mediated focussing allows autorecognition to be physiological, not pathological: a large proportion of patent repertoire TCR clonotypes can have autoantigens among their potential antigens, but autoimmunity is

averted provided the CD8-modulatory system keeps an activated clonotype "trained" on the salient epitope.

The present model postulates the existence of favourable membrane microdomains, to which TCRs can be recruited only when bound to pMHC I. Such domains have been proposed to exist in the form of lipid rafts, membrane areas enriched in cholesterol and glycosphingolipids which serve as the sites of colocalization of TCR/CD3, protein tyrosine kinases, and adaptor molecules (Arcaro et al., 2001; Montixi et al., 1998; Bosselut et al., 1999; Janes et al., 1999; Brdičková et al., 2003). Partitioning of CD8 to lipid rafts is crucially dependent on palmitoylation of the cytoplasmic tail of CD8 β (Arcaro et al., 2000, 2001). Therefore, inasmuch as the increased rate of phosphorylation is dependent on recruitment to microdomains (rather than direct recruitment of p56^{lck} by CD8 to the TCR/CD3 complex, which CD8 $\alpha\alpha$ could mediate), the main action of CD8 $\alpha\alpha$ would be to modulate on- and off-rates. Either type of dimer is able to reduce the TCR/pMHC I dissociation rate since CD8 α binds MHC I, (Kern et al., 1998; Wyer et al., 1999). The contrast between points f and g in Fig. 2 illustrates the opposite effects which may be exerted by upregulation of CD8 $\alpha\alpha$, which only affects the mean TCR/pMHC I dwell time and CD8 $\alpha\beta$ which affects lipid raft colocalization (cf. Cawthon and Alexander-Miller, 2002; Gangadharan and Cheroutre, 2004). Thus, a T cell may be able to shift its focus to a distinct subset of its potential agonists by altering the relative expression levels of CD8 $\alpha\alpha$ and CD8 $\alpha\beta$.

The off-rate measured with HLA DT227/8KA depends only on v^{fast} , whereas the off-rate found with the wild-type HLA is a mixture of both v^{fast} and v^{slow} , as explained above in the section on mean dwell-time modulation. Thus, the effective dwell-time prolongation of ~ 1.5 as measured with hTERT_{540–548} APLs is a lower bound to the actual CD8-mediated stabilization factor v^{fast}/v^{slow} . Analysis based on HLA A2 mutants with a range of pMHC I/CD8 affinities indicated that stabilization can exceed a factor 2 (Wooldridge et al., 2005; Laugel et al., 2007); in physiological terms, this factor corresponds to the extremes of CD8 expression, i.e. from no expression to MHC I-saturating levels. Similar remarks apply to the enhancement of the on-rate. Thus, the data presented here support the claim that the pMHC I/CD8 interaction prolongs dwell time by at least a factor 1.5, and also enhances the on-rate by at least a factor 1.5, with the important caveat that this latter figure is derived from two model estimates (the affinity effect in the equilibrium staining curve and the off-rate effect), so that the strength of the on-rate effect may well be much greater or smaller.

Moreover, the on-rate effect may differ among APL variants. The empirical rate λ^{fast} in the association experiments estimates how rapidly $\lambda_{eff}(t)$ relaxes to its equilibrium value. The rate which may be expected to dominate the initial rapid phase is $\partial_1 R_T + v$; indeed, APL variants 3G and 8Y show that λ^{fast} broadly agrees with the off-rate v , and λ^{fast} increases upon abrogation of the

pMHC/CD8 interaction, as expected when the rate is dominated by v . For variant 8T, however, the initial rate λ^{fast} is much faster, and abrogation of the pMHC/CD8 interaction acts to *decrease* this rate somewhat; again this points to a greater role of on-rate effects, as opposed to off-rate effects, in coreceptor modulation for this intermediate affinity ligand.

The dominant (slow) rate in the biphasic association curves arises from an intricate interplay between the various bound forms depicted in Fig. 1. To interpret the rate estimates listed in the final column of Table 5, we must also consider the estimated proportions of triplets, Table 3, and Eq. (14) for the effective rate. The apparently much more profound effect on APL variant 3G can be understood from the fact that λ_{eff} is determined almost exclusively by $3\psi + v_3$, since the staining is almost entirely in the form of triplets. Since v_3 is relatively small, the decrease in ψ due to abrogation of the pMHC/CD8 interaction (“capture effect”) is the dominant effect on the effective rate of association. For variants 8Y and 8T, the duplet proportions are higher. In the duplet factor, the decrease in the term 2ψ is partially compensated for by the increase in v_2 . This explains why, for 8Y and 8T, the final two columns of Table 5 show a far less marked reduction of $\lambda_{eff}(\infty)$ when the MHC/CD8 interaction is abrogated.

A second indirect piece of evidence pointing to TCR/pMHC on-rate modulation by CD8 is the marked effect of the abrogation of the pMHC/CD8 interaction on MHC tetramer capture (Table 5). The pronounced capture effect we report is consistent with the analysis of Gakamsky et al. (2005), who suggest that CD8 is likely to play a major role in capturing a pMHC tetramer from solution. Our analysis points to roughly equal contributions of on-rate and off-rate enhancement on affinity modulation by CD8. However, if the rate of tetramer capture is closely linked to the single-site on-rate in the cell:cell contact environment, the contribution of the on-rate might be much more important, particularly in ligands with relatively high TCR/pMHC dissociation constants.

In terms of the collision theory of molecular reaction dynamics, the effect of CD8 on the TCR/pMHC association rate may be viewed as a modulation of either the activation energy or the reactive cross-section, or both, although the former seems less likely since the interaction between CD8 and MHC is relatively weak (Purbhoo et al., 2001). Modulation of the reactive cross-section can be understood as an increase of the steric factor characterizing TCR/pMHC collisions, with the CD8 molecule effectively acting as a grappling hook; this picture would be consistent with the finding that favourable entropic forces rather than enthalpic forces play a major role in (at least some) TCR/pMHC interactions (Ely et al., 2006). Indirect evidence supporting the notion of CD8 promoting the formation of TCR/pMHC contacts, even with non-cognate peptide presented by MHC, has recently been put forward by Anikeeva et al. (2006).

The physiological significance of TCR/pMHC on-rate modulation is rather subtle. The formula $v \exp\{-vT_R\}$ for the rate of productive ligand engagement suggests that the single-site off-rate v and the receptor triggering threshold T_R are the sole determinants of T cell activation. However, this formula applies only under MHC-limited conditions (van den Berg et al., 2002); the full formula is $v \exp\{-vT_R\}(1 + K_D^{TCR/pMHC}/[R])^{-1}$ where $[R]$ is the surface density of free TCRs. This full formula reduces to the simpler (MHC-limited) expression provided that $[R] \gg K_D^{TCR/pMHC}$. On-rate enhancement, which leads to a smaller $K_D^{TCR/pMHC}$, allows this condition to be satisfied at lower TCR-expression levels. The curves in Fig. 2 indicate that there is an optimal TCR/pMHC off-rate; the theory of van den Berg et al. (2002) predicts that the optimum broadens in the TCR-limited regime, as has recently been confirmed experimentally (Gonzales et al., 2005). Thus, the physiological significance of on-rate modulation (i.e. improvement of affinity through reduction of the TCR/pMHC dissociation constant) may be to maintain the sharp optimum in the dependence of triggering on off-rate. This effect would be more important for ligands of intermediate to low affinity.

The fast and slow rates of MHC-tetramer staining kinetics contain information of duplet/triplet ratios in tetramer/TCR associations. Since tetramers are artificial constructs, such data may seem to have no direct physiological significance. However, therapeutic use of MHC tetramers has been proposed (Sakita et al., 1996; Maile et al., 2001). MHC oligomers are capable of inducing T cell activation, indicating that TCR clusters are competent nucleation points for signalosome formation; experiments with MHCII monomers, dimers, trimers, and tetramers indicate that both duplet and triplet clusters are competent (Boniface et al., 1998; Cochran et al., 2000). If duplets and triplets are not equally potent, an experimental means of gauging triplet/duplet ratios would be of clinical value.

In summary, we have used pMHC tetramer-based kinetic analysis, combined with target cells expressing MHC mutants, to dissect the various modulatory effects through which the coreceptor CD8 differentially modulates TCR sensitivity to its various potential agonists, and thereby modulates TCR specificity in a dynamic, tunable fashion. Taken together, the experimental findings suggest that CD8 has an influence on TCR/pMHC affinity through modulation of both the on-rate and the off-rate. Furthermore, we have presented evidence that CD8 modulates the TCR triggering threshold by recruiting TCRs to favourable membrane microdomains in which progression to signalling-competent signalosomes proceeds much faster. These effects endow the T cell with an exquisite means of tuning its TCR to the pertinent peptide ligand by adjusting the expression levels of CD8 $\alpha\alpha$ and/or CD8 $\alpha\beta$. We suggest that at the level of the whole adaptive immune system, CD8-based modulation of TCR specificity

vastly amplifies the functional diversity of the finite number of TCR clonotypes present in the T cell repertoire.

Acknowledgements

We are grateful to an anonymous referee for comments which helped improve this paper.

References

- Alexander-Miller, M.A., 2000. Differential expansion and survival of high and low avidity cytotoxic T cell populations during the immune response to a viral infection. *Cell. Immunol.* 201, 58–62.
- Anderton, S.M., Wraith, D.C., 2002. Selection and fine-tuning of the autoimmune T cell repertoire. *Nat. Rev. Immunol.* 2, 487–497.
- Anikeeva, N., Lebedeva, T., Clapp, A.R., Goldman, E.R., Dustin, M.L., Mattoussi, H., Sykulev, Y., 2006. Quantum dot/peptide-MHC biosensors reveal strong CD8-dependent cooperation between self and viral antigens that augment the T cell response. *Proc. Natl Acad. Sci. USA* 103, 16846–16851.
- Arcaro, A., Grégoire, C., Boucheron, N., Stolz, S., Palmer, E., Malissen, B., Luescher, I.F., 2000. Essential role of CD8 palmitoylation in CD8 coreceptor function. *J. Immunol.* 165, 2068–2076.
- Arcaro, A., Grégoire, C., Bakker, T.R., Baldi, L., Jordan, M., Goffin, L., Boucheron, N., Wurm, F., van der Merwe, P.A., Malissen, B., Luescher, I.F., 2001. CD8 β endows CD8 with efficient coreceptor function by coupling T cell receptor/CD3 to raft-associated CD8/p56^{lck} complexes. *J. Exp. Med.* 194, 1485–1495.
- Ashton-Rickardt, P.G., Tonegawa, S., 1994. A differential-avidity model for T-cell selection. *Immunol. Today* 15, 362–366.
- Bankovich, A.J., Girvin, A.T., Moesta, A.K., Garcia, K.C., 2004. Peptide register shifting within the MHC groove: theory becomes reality. *Mol. Immunol.* 40, 1033–1039.
- van den Berg, H.A., Molina-París, C., 2003. Thymic presentation of autoantigens and the efficiency of negative selection. *J. Theor. Med.* 5, 1–22.
- van den Berg, H.A., Rand, D.A., 2004a. Dynamics of T cell activation threshold tuning. *J. Theor. Biol.* 228, 397–416.
- van den Berg, H.A., Rand, D.A., 2004b. Foreignness as a matter of degree: the relative immunogenicity of peptide/MHC ligands. *J. Theor. Biol.* 231, 535–548.
- van den Berg, H.A., Rand, D.A., 2007. Quantitative theories of T-cell responsiveness. *Immunol. Rev.* 216, 81–92.
- van den Berg, H.A., Rand, D.A., Burroughs, N.J., 2001. A reliable and safe T cell repertoire based on low-affinity T cell receptors. *J. Theor. Biol.* 209, 465–486.
- van den Berg, H.A., Burroughs, N.J., Rand, D.A., 2002. Quantifying the strength of ligand antagonism in TCR triggering. *Bull. Math. Biol.* 64, 781–808.
- Blok, R., Margulies, D.H., Pease, L., Ribaud, R.K., Schneck, J., McCluskey, J., 1996. CD8 expression alters the fine specificity of an alloreactive MHC class I-specific T hybridoma. *Int. Immunol.* 4, 455–466.
- Bongrand, P., Malissen, B., 1998. Quantitative aspects of T-cell recognition: from within the antigen-presenting cell to within the T cell. *BioEssays* 20, 412–422.
- Boniface, J.J., Rabinowitz, J.D., Wülfing, C., Hampl, J., Reich, Z., Altman, J.D., Kantor, R.M., Beeson, C., McConnell, H.M., Davis, M.M., 1998. Initiation of signal transduction through T cell receptor requires the peptide multivalent engagement of MHC ligands. *Immunity* 9, 459–466.
- Bosselut, R., Zhang, W., Ashe, J.M., Kopacz, J.L., Samuelson, L.E., Singer, A., 1999. Association of the adaptor molecule LAT with CD4 and CD8 coreceptors identifies a new coreceptor function in T cell receptor signal transduction. *J. Exp. Med.* 190, 1517–1525.
- Bosselut, R., Kubo, S., Guinter, T., Kopacz, J.L., Altman, J.D., Feigenbaum, L., Singer, A., 2000. Role of CD8 β domains in CD8 coreceptor function: importance for MHC I binding, signaling, and positive selection of CD8⁺ T cells in the Thymus. *Immunity* 12, 409–418.
- Brdičková, N., Brdička, T., Angelisová, P., Horváth, O., Špiča, J., Hilgert, I., Pačes, J., Simeoni, L., Kliche, S., Merten, C., Schraven, B., Hofeš, V., 2003. LIME: a new membrane raft-associated adaptor protein involved in CD4 and CD8 coreceptor signaling. *J. Exp. Med.* 198, 1453–1462.
- Bunnell, S.C., Hong, D.I., Kardon, J.R., Yamazaki, T., McGlade, C.J., Barr, V.A., Samelson, L.E., 2002. T cell receptor ligation induces the formation of dynamically regulated signaling assemblies. *J. Cell Biol.* 158, 1263–1275.
- Burroughs, N., Wedagedera, J., 2006. T cell activation: a queueing theory analysis at low agonist density. *Biophys. J.* 91, 1604–1618.
- Cawthon, A.G., Alexander-Miller, M.A., 2002. Optimal colocalization of TCR and CD8 as a novel mechanism for the control of functional avidity. *J. Immunol.* 169, 3492–3498.
- Chan, C., George, A.J.T., Stark, J., 2003. T cell sensitivity and specificity—kinetic proofreading revisited. *Discrete and Continuous Dynamical Syst. Ser. B* 3, 343–360.
- Cochran, J.R., Cameron, T.O., Stern, L.J., 2000. The relationship of MHC-peptide binding and T cell activation probed using chemically defined MHC class II oligomers. *Immunity* 12, 241–250.
- Daniels, M.A., Devine, L., Miller, J.D., Moser, J.M., Lukacher, A.E., Altman, J.D., Kavathas, P., Hogquist, K.A., Jameson, S., 2001. CD8 binding to MHC class I molecules is influenced by T cell maturation and glycosylation. *Immunity* 15, 1051–1061.
- Davis, S.J., van der Merwe, P.A., 2006. The kinetic-segregation model: TCR triggering and beyond. *Nat. Immunol.* 7, 803–809.
- Doucey, M.-A., Legler, D., Boucheron, N., Cerottini, J.-C., Bron, C., Luescher, I.F., 2001. CTL activation is induced by cross-linking of TCR/MHC-peptide-CD8/p56^{lck} adducts in rafts. *Eur. J. Immunol.* 31, 1561–1570.
- Dunbar, P.R., Ogg, G.S., Chen, J., Rust, N., van der Bruggen, P., Cerundolo, V., 1998. Direct isolation, phenotyping and cloning of low-frequency antigen-specific cytotoxic T lymphocytes from peripheral blood. *Curr. Biol.* 8, 413–662.
- Ely, L.K., Beddoe, T., Clements, C.S., Matthews, J.M., Purcell, A.W., Kjer-Nielsen, L., McCluskey, J., Rossjohn, J., 2006. Disparate thermodynamics governing T cell receptor–MHC-I interactions implicate extrinsic factors in guiding MHC restriction. *Proc. Natl Acad. Sci. USA* 103, 6641–6646.
- Filipp, D., Zhang, J., Leung, B.L., Shaw, A., Levin, S.D., Veillette, A., Julius, M., 2003. Regulation of Fyn through translocation of activated Lck into lipid rafts. *J. Exp. Med.* 197, 1221–1227.
- Gakamsky, D.M., Luescher, I.F., Pramanik, A., Kopito, R.B., Lemonnier, F., Vogel, H., Rigler, R., Pecht, I., 2005. CD8 kinetically promotes ligand binding to the T-cell antigen receptor. *Biophys. J.* 89, 2121–2133.
- Gangadharan, D., Cheroutre, H., 2004. The CD8 isoform CD8 $\alpha\alpha$ is not a functional homologue of the TCR co-receptor CD8 $\alpha\beta$. *Curr. Opin. Immunol.* 16, 264–270.
- Gavin, M.A., Bevan, M.J., 1995. Increased peptide promiscuity provides a rationale for the lack of N regions in the neonatal T cell repertoire. *Immunity* 3, 793–800.
- Goldrath, A.W., Bevan, M.J., 1999. Selecting and maintaining a diverse T-cell repertoire. *Nature* 402, 255–262.
- Gonzales, P.A., Carreño, L.J., Coombs, D., Mora, J.E., Palmeiri, E., Goldstein, B., Nathanson, S.G., Kalergis, A.M., 2005. T cell receptor binding kinetics required for T cell activation depend on the density of the cognate ligand of the antigen-presenting cell. *Proc. Natl Acad. Sci. USA* 102, 4824–4829.
- Gross, D.-A., Graff-Dubois, S., Opolon, P., Cornet, S., Alves, P., Bencecurel-Griscelli, A., Faure, O., Guillaume, P., Firat, H., Chouaib, S., Lemonnier, F.A., Davoust, J., Miconnet, I., Vonderheide, R.H., Kosmatopoulos, K., 2004. High vaccination efficiency of low-affinity epitopes in antitumor immunotherapy. *J. Clin. Invest.* 113, 425–433.

- Hemmer, B., Stefanova, I., Vergelli, M., Germain, R.N., Martin, R., 1998. Relationships among TCR ligand potency, thresholds for effector function elicitation, and the quality of early signaling events in human T cells. *J. Immunol.* 160, 5807–5814.
- Holler, P.D., Kranz, D.M., 2003. Quantitative analysis of the contribution of TCR/pepMHC affinity and CD8 to T cell activation. *Immunity* 18, 255–264.
- Holler, P.D., Kranz, D.M., 2004. T cell receptors: affinities, cross-reactivities, and a conformer model. *Mol. Immunol.* 40, 1027–1031.
- Holler, P.D., Chlewicki, L.K., Kranz, D.M., 2002. TCRs with high affinity for foreign pMHC show self-reactivity. *Nat. Immunol.* 4, 55–62.
- Hudrisier, D., Kessler, B., Valitutti, S., Horvath, C., Cerottini, J.-C., Luescher, I.F., 1998. The efficiency of antigen recognition by CD8⁺ CTL clones is determined by the frequency of serial TCR engagement. *J. Immunol.* 161, 553–562.
- Hutchinson, S.L., Wooldridge, L., Tafuro, S., Laugel, B., Glick, M., Boulter, J.M., Jakobsen, B.K., Price, D.A., Sewell, A.K., 2003. The CD8 T cell coreceptor exhibits disproportionate biological activity at extremely low binding affinities. *J. Biol. Chem.* 278, 24285–24293.
- Ignatowicz, L., Kappler, J., Marrack, P., 1996. The repertoire of T cells shaped by a single MHC/peptide ligand. *Cell* 84, 521–529.
- Janes, P.W., Ley, S.C., Magee, A.I., 1999. Aggregation of lipid rafts accompanies signalling via the T cell antigen receptor. *J. Cell Biol.* 147, 447–461.
- Janeway Jr., C.A., 1992. The T cell receptor as a multicomponent signalling machine: CD4/CD8 coreceptors and CD45 in T cell activation. *Annu. Rev. Immunol.* 10, 645–674.
- Kalergis, A.M., Boucheron, N., Doucey, M.-A., Palmieri, E., Goyarts, E.C., Vegh, Z., Luescher, I.F., Nathanson, S.G., 2001. Efficient T cell activation requires an optimal dwell-time of interaction between the TCR and the pMHC complex. *Nat. Immunol.* 2, 229–234.
- Kern, P.S., Teng, M.K., Smolyar, A., Liu, J.H., Liu, J., Hussey, R.E., Spoerl, R., Chang, H.C., Reinherz, E.L., Wang, J.H., 1998. Structural basis of CD8 coreceptor function revealed by crystallographic analysis of a murine CD8 $\alpha\alpha$ ectodomain fragment in complex with H-2Kb. *Immunity* 9, 519–530.
- Kersh, G.J., Allen, P.M., 1996. Structural basis for T cell recognition of altered peptide ligands: a single T cell receptor can productively recognize a large continuum of related ligands. *J. Exp. Med.* 184, 1259–1268.
- Laugel, B., Boulter, J.M., Lissin, N., Vuidepot, A., Li, Y., Gostick, E., Crotty, L.E., Douek, D.C., Hemelaar, J., Price, D.A., Jakobsen, B.K., Sewell, A.K., 2005. Design of soluble recombinant T cell receptors for antigen targeting and T cell inhibition. *J. Biol. Chem.* 280, 1882–1892.
- Laugel, B., van den Berg, H.A., Gostick, E., Cole, D.K., Wooldridge, L., Boulter, J., Milicic, A., Price, D.A., Sewell, A.K., 2007. Different T cell receptor affinity thresholds and CD8 coreceptor dependency govern cytotoxic T lymphocyte activation and tetramer binding properties. *J. Biol. Chem.* 282, 23799–23810.
- Luescher, I.F., Vivier, E., Layer, A., Mahiou, J., Godeau, F., Malissen, B., Romero, P., 1995. CD8 modulation of T-cell antigen receptor–ligand interactions on living cytotoxic T lymphocytes. *Nature* 373, 353–356.
- Lyons, D.S., Lieberman, S.A., Hampl, J., Boniface, J., Chien, Y.-h., Berg, L.J., Davis, M.M., 1996. A TCR binds to antagonist ligands with lower affinities and faster dissociation rates than to agonists. *Immunity* 5, 53–61.
- Lyons, G.E., Moore, T., Brasic, N., Li, M., Roszkowski, J.J., Nishimura, M.I., 2006. Influence of human CD8 on antigen recognition by T-cell receptor-transduced cells. *Cancer Res.* 66, 11455–11461.
- Maile, R., Wang, B., Schooler, W., Meyer, A., Collins, E.J., Frelinger, J.A., 2001. Antigen-specific modulation of an immune response by *in vivo* administration of soluble MHC class I tetramers. *J. Immunol.* 167, 3708–3714.
- Maile, R., Siler, C.A., Kerry, S.E., Midkiff, K.E., Collins, E.J., Frelinger, J.A., 2005. Peripheral “CD8 tuning” dynamically modulates the size and responsiveness of an antigen-specific T cell pool *in vivo*. *J. Immunol.* 174, 619–627.
- Malissen, B., 2003. An evolutionary and structural perspective on T cell antigen receptor function. *Immunol. Rev.* 191, 7–27.
- Mason, D., 1998. A very high level of crossreactivity is an essential feature of the T-cell receptor. *Immunol. Today* 19, 395–404.
- Mason, D., 2001. Some quantitative aspects of T cell repertoire selection: the requirement for regulatory T cells. *Immunol. Rev.* 182, 80–88.
- Matsui, K., Boniface, J.J., Steffner, P., Reay, P.A., Davis, M.M., 1994. Kinetics of T-cell receptor binding to peptide/I-E^k complexes: correlation of the dissociation rate with T-cell responsiveness. *Proc. Natl Acad. Sci. USA* 91, 12862–12866.
- Montixi, C., Langlet, C., Bernard, A.-M., Thimonier, J., Dubois, C., Wurbel, M.-A., Chauvin, J.-P., Pierres, M., He, H.-T., 1998. Engagement of T cell receptor triggers its recruitment to low-density membrane domains. *EMBO J.* 17, 5334–5348.
- Nicholson, L.B., Anderson, A.C., Kuchroo, V.K., 2000. Tuning T cell activation threshold and effector function with cross-reactive peptide ligands. *Int. Immunol.* 12, 205–213.
- Pecht, I., Gakamsky, D.M., 2005. Spatial coordination of CD8 and TCR molecules controls antigen recognition by CD8⁺ T-cells. *FEBS Lett.* 579, 3336–3341.
- Purbhoo, M.A., Boulter, J.M., Price, D.A., Vuidepot, A.-L., Hourigan, C.S., Dunbar, P.R., Olson, K., Dawson, S.J., Phillips, R.E., Jakobsen, B.K., Bell, J.I., Sewell, A.K., 2001. The human CD8 coreceptor effects cytotoxic T cell activation and antigen sensitivity primarily by mediating complete phosphorylation of the T cell receptor ζ chain. *J. Biol. Chem.* 276, 32786–32792.
- Sakita, I., Högri, H., Rui, S., Fuming, W., Nathanson, S.G., 1996. *In vivo* CTL immunity can be elicited by *in vitro* reconstituted MHC/peptide complex. *J. Immunol. Meth.* 192, 105–115.
- Schade, A.E., Levine, A.D., 2003. Phosphatases in concert with kinases set the gain for signal transduction through the T cell receptor. *Mol. Immunol.* 40, 531–537.
- Seddon, B., Mason, D., 2000. The third function of the thymus. *Immunol. Today* 21, 95–99.
- Slifka, M.K., Whitton, J.L., 2001. Functional avidity maturation of CD8⁺ T cells without selection of higher affinity TCR. *Nat. Immunol.* 2, 711–717.
- Valitutti, S., Lanzavecchia, A., 1997. Serial triggering of TCRs: a basis for the sensitivity and specificity of antigen recognition. *Immunol. Today* 18, 299–304.
- Viola, A., Lanzavecchia, A., 1996. T cell activation determined by T cell receptor number and tunable thresholds. *Science* 273, 104–106.
- Wilson, D.B., Wilson, D.H., Schroder, K., Pinilla, C., Blondelle, S., Houghten, R.A., Garcia, K.C., 2004. Specificity and degeneracy of T cells. *Mol. Immunol.* 40, 1047–1055.
- Wooldridge, L., van den Berg, H.A., Glick, M., Gostick, E., Brenchley, J.M., Douek, D.C., Price, D.A., Sewell, A.K., 2005. Interaction between the CD8 coreceptor and MHC class I stabilizes TCR–antigen complexes at the cell surface. *J. Biol. Chem.* 280, 27491–27501.
- Wyer, J.R., Willcox, B.E., Gao, G.F., Gerth, U.C., Davis, S.J., Bell, J.I., van der Merwe, P.A., Jakobsen, B.K., 1999. T cell receptor and co-receptor CD8 $\alpha\alpha$ bind peptide–MHC independently and with distinct kinetics. *Immunity* 10, 219–225.